

PRODUCTION OF OPTICALLY ACTIVE ALCOHOL BY REDUCTION WITH ASYMMETRIC BORANE

Patent Number: JP9255592

Publication date: 1997-09-30

Inventor(s): MASUI MORIYASU

Applicant(s):: SHIONOGI & CO LTD

Requested
Patent: ☐ JP9255592Application
Number: JP19960246276 19960918Priority Number
(s):IPC Classification: C07B53/00 ; B01J31/14 ; C07B41/02 ; C07B43/04 ; C07C29/143 ; C07C31/125 ; C07C33/22 ;
C07C213/00 ; C07C215/10 ; C07C215/44 ; C07C235/04 ; C07C251/36EC
Classification:


Equivalents:

Abstract


PROBLEM TO BE SOLVED: To obtain an optically active alcohol which is useful as a medicine, agrochemical or the like in high asymmetric yield by asymmetric reduction of an unsymmetrical ketone with hydroborane in the presence of a chiral β -aminoalcohol and a trialkoxyborane.

SOLUTION: (A) A compound formula I ($R<1>$ and $R<2>$ are each an organic residue different from each other) is reduced with (D) hydroborane in the presence of (B) a chiral β -aminoalcohol of formula II [$R<3>$ is an alkyl, a cycloalkyl, a (substituted) aryl; $R<4>$ and $R<5>$ are each H, an alkyl; $R<6>$ is H, an alkyl, a cycloalkyl; * is an asymmetric carbon] and (C) a compound of formula III ($R<7>$ is an alkyl) to give an optically active alcohol of formula IV ($R<1>$ and $R<2>$ are each an organic residue same as $R<1>$ and $R<2>$ respectively). The component A is preferably, for example, acetophenone, propiophenone, phenacyl chloride, benzyl phenyl ketone. An α -oxoketoxime derivative also may be used as a component A.

Data supplied from the esp@cenet database - I2




I



II



III



IV